

Fast
action



Rumilab®

Dietary feed supplement for veterinary use



Indicated for indigestion

- ✓ Reduces the risk of acidosis
- ✓ Reduces the risk of ketosis
- ✓ Indicated for ruminal atony
- ✓ Treatment against anorexia

Faster action

- ✓ High concentration per intake
- ✓ Alternative source of B group vitamins and minerals
- ✓ Provides raw material for the maintenance of the adequate ruminal flora

Convenience of use


- ✓ Package with two sachets
- ✓ One treatment for each animal
- ✓ Maximum efficacy in treatments combined with Rumintral

LABIANA

always works



Development of the clinical picture of acidosis in ruminants



COMPOSITION

Glycerol (4.64 g/kg), brewer's yeast (dried), casein powder, dicalcium phosphate, sodium chloride.

Additives (per 175 g bag):

Vitamins, provitamins and chemically well-defined substances of similar effect:

Vitamin B2/ Riboflavin.....	50 mg
Niacinamide (3a315)	25 mg
Vitamin B1/Thiamine chlorhydrate (3a820)	15 mg

Amino acids, their salts and similar:

DL-Methionine, technically pure (3c301)	3 g
---	-----

Trace elements or compounds of trace elements:

Manganese (E5) (manganese oxide)	200.26 g
Zinc (E6) (heptahydrate zinc sulfate)	2.27 g
Iron (E1) (Ferrous carbonate)	1.04 g

Preservatives:

Sodium propionate (E281)	45 g
Calcium propionate (E282)	30 g

INDICATIONS

Dairy cows and calves

- Reduction of the risk of acidosis, recommended period of use: maximum 2 months.
- Especially indicated for high-yield dairy cows and for intensive fattening calves.
- Reduction of risk of ketosis in dairy cows.
- It is also an alternative source of B group vitamins and minerals.

DOSAGE AND ROUTE OF ADMINISTRATION

Reduction of the risk of ketosis:

Dairy cows:

1 bag (175 g), 1 or 2 times per day or 1/2 bag (87.5 g) / day, for 2-3 days.

Reduction of the risk of acidosis:

Dairy cows: 1 bag (175 g) 1 or 2 times/day or 1/2 bag (87.5 g)/day for 2-3 days.

Calves: 1/2 bag (87.5 g) 1 or 2 times/day or 1/4 bag (43.75 g)/day for 2-3 days.

Add the indicated doses at 1 kg of food to one bag of mixture, in 1/2 kg of food at other dosages.

Oral route, in the food.

Request the opinion of a veterinary before using it or before extending the period of use.

WITHDRAWAL PERIOD

Not required.

USE DURING PREGNANCY

Dairy cows: 3 to 6 weeks after calving.

PRESENTATION

Box with 2 sachets of 175 g each.

αESP-08100341